

INTERACTING NONCOMMUTATIVE LUMPS

CORNELIU SOCHICHIU

*Bogoliubov Laboratory of Theoretical Physics
Joint Institute for Nuclear Research*

141980 Dubna, Moscow Reg.

RUSSIA

Institutul de Fizică Aplicată AȘ,

str. Academiei, nr. 5,

MD2028 Chișinău,

MOLDOVA

E-mail: sochichi@thsun1.jinr.ru

We consider interaction of two lumps corresponding to 0-branes in noncommutative gauge theory

Recent development of string theory has shown that noncommutative (NC) models can arise in certain limits of String Theory ¹.

Noncommutative field theory operates with operators defined on a Hilbert space rather than with ordinary functions. There is, however, a one-to-one correspondence between such operators and functions which is given by the Weyl map. Important solutions discovered recently are the so called noncommutative solitons or lumps ². These configurations are represented by projector operators. Their Weyl symbols are localised functions.

In this short note we are going to consider the interaction of two such gauge field lumps. This is realised by the dynamics of the configuration which represents a superposition of two projectors to finite-dimensional subspaces of the Hilbert space which correspond to two copies of the oscillator vacuum state shifted along noncommutative plane. For more details the reader is referred to Ref. ³.

0.1 The Model.

Consider the noncommutative gauge model described by the following action,

$$S = \int dt \operatorname{tr} \left(\frac{1}{2} \dot{X}^i \dot{X}^i + \frac{1}{4g^2} [X^i, X^j]^2 \right). \quad (1)$$

Here fields X^i , $i = 1, \dots, D$ are time dependent Hermitian operators defined on some separable infinite-dimensional Hilbert space $\mathcal{H}$.

In the two-dimensional Moyal–Weyl form some operators f are represented by functions $f(x)$ on x^μ $\mu = 1, 2$, subject to the star-product,

$$f * g(x) = e^{i\theta \epsilon^{\mu\nu} \partial_\mu \partial'_\nu} f(x) g(x') \Big|_{x'=x}, \quad (2)$$

where x^μ satisfy, $[x^1, x^2] = i\theta$.

Equations of motion look as follows

$$\ddot{X}_i + \frac{1}{g^2} [X_i, [X_i, X_j]] = 0. \quad (3)$$

In the case of just two lumps one can take them to be:

$$X_1 = cVPV^{-1} \equiv c|-u/2\rangle \langle -u/2|, \quad (4a)$$

$$X_2 = cV^{-1}PV \equiv c|u/2\rangle \langle u/2| \quad (4b)$$

$$X_i = \text{const}, \quad i = 3, \dots, D, \quad (4c)$$

where we introduced the shorthand notations,

$$V = e^{(i/2)p_\mu u^\mu} = e^{\frac{1}{2}(a\bar{u} - \bar{a}u)}, \quad (5)$$

$$P = |0\rangle \langle 0|. \quad (6)$$

To find the time evolution of such a system one has to solve the equations of motion (3) with initial condition given by (4) and $\dot{X}_i|_{t=0} = 0$.

Symmetries of the model and initial conditions allows one to reduce the system to a two-dimensional particle described by the equations,

$$\ddot{X} = -\frac{2}{g^2} Y^2 X, \quad (7a)$$

$$\ddot{Y} = -\frac{2}{g^2} X^2 Y, \quad (7b)$$

with the initial conditions,

$$X|_{t=0} = e^{-\frac{1}{2}|u|^2}, \quad Y|_{t=0} = -\sqrt{1 - e^{-|u|^2}}. \quad (7c)$$

This model has a number of interesting features^{4,5,6}. The motion resulting from above equations appears to be *stochastic* for all values of u except $u = \sqrt{\theta} \ln 2$. When $u = \sqrt{\theta} \ln 2$ the motion is periodic but unstable, infinitesimal deviation from this position brings it back to stochasticity.

The solution to equations (3) is expressed in terms linear combination of some functions $\sigma_a(\bar{z}, z)$ with coefficients depending on $X(t)$ and $Y(t)$,

$$X_1(t, \bar{z}, z) = \frac{1}{2} \sigma_0(\bar{z}, z) + X(t) \sigma_3(\bar{z}, z) + Y(t) \sigma_1(\bar{z}, z), \quad (8)$$

$$X_2(t, \bar{z}, z) = \frac{1}{2} \sigma_0(\bar{z}, z) + X(t) \sigma_3(\bar{z}, z) - Y(t) \sigma_1(\bar{z}, z). \quad (9)$$

Functions $\sigma_a(\bar{z}, z)$ are the Weyl symbols of the two-dimensional Pauli matrices,

$$\sigma_1(z, \bar{z}) = \frac{2}{\sqrt{1 - e^{-|u|^2}}} \left(e^{-2|z - \frac{u}{2}|^2} - e^{-2|z + \frac{u}{2}|^2} \right), \quad (10a)$$

$$\sigma_2(z, \bar{z}) = \frac{2ie^{-2\bar{z}z}}{\sqrt{1 - e^{-|u|^2}}} \left(e^{\bar{z}u - \bar{u}z} - e^{-\bar{z}u + \bar{u}z} \right), \quad (10b)$$

$$\begin{aligned} \sigma_3(z, \bar{z}) = & -\frac{2e^{-\frac{1}{2}|u|^2}}{1 - e^{-|u|^2}} \left(e^{-2|z - \frac{u}{2}|^2} + e^{-2|z + \frac{u}{2}|^2} \right) \\ & + \frac{e^{-2\bar{z}z}}{1 - e^{-|u|^2}} \left(e^{\bar{z}u - \bar{u}z} + e^{-\bar{z}u + \bar{u}z} \right), \end{aligned} \quad (10c)$$

$$\begin{aligned} \mathbb{I}(\bar{z}, z) = \sigma_0(z, \bar{z}) = & \frac{2}{1 - e^{-|u|^2}} \left(e^{-2|z - \frac{u}{2}|^2} + e^{-2|z + \frac{u}{2}|^2} \right) \\ & - \frac{2e^{-2|z|^2 - \frac{1}{2}|u|^2}}{1 - e^{-|u|^2}} \left(e^{\bar{z}u - \bar{u}z} + e^{-\bar{z}u + \bar{u}z} \right). \end{aligned} \quad (10d)$$

The solution describes lumps bouncing (stochastically) around “points” $z = 0, \pm u/2$ of the noncommutative plane. In the string theory language the heights of the lumps can be interpreted as transversal (to the noncommutative 2-brane) coordinates of two 0-branes.

Acknowledgments

This work was supported by RFBR grant #99-01-00190, INTAS grants #00 262, Scientific School support grant # 00-15-96046.

References

1. N. Seiberg and E. Witten, *JHEP* **09** (1999) 032, [hep-th/9908142](#).
2. R. Gopakumar, S. Minwalla, and A. Strominger, *JHEP* **05** (2000) 020, [hep-th/0003160](#).
3. C. Sochichiu, [hep-th/0104076](#).
4. G. Z. Baseian, S. G. Matinian, and G. K. Savvidy, *Pisma Zh. Eksp. Teor. Fiz.* **29** (1979) 641.
5. B. V. Medvedev, *Theor. Math. Phys.* **60** (1985) 782–797.
6. I. Y. Aref’eva, A. S. Koshelev, and P. B. Medvedev, in matrix theory,” *Mod. Phys. Lett.* **A13** (1998) 2481, [hep-th/9804021](#).